

Neural Correlates of Self-Focused and Other-Focused Strategies for Coping With Cigarette Cue Exposure

Stephen J. Wilson
The Pennsylvania State University

Michael A. Sayette
University of Pittsburgh

Julie A. Fiez
University of Pittsburgh and Center for the Neural Basis of Cognition

Brain imaging research has begun to characterize the neurocognitive processes that cigarette smokers utilize to cope with cue-elicited craving. Presently, however, it remains unclear whether distinct neural substrates support different types of coping. We sought to address this knowledge gap by examining neural responses associated with self-focused and other-focused coping techniques. Fifty-seven treatment-seeking male cigarette smokers initiated an attempt to quit smoking and subsequently underwent functional MRI, during which they were asked to hold and view neutral cues and a cigarette. Participants were instructed to engage in either self-focused or other-focused coping while being presented with the cigarette and an opportunity to smoke. Those who were told to engage in self-focused coping, but not those told to utilize other-focused coping, exhibited significant activation of several regions previously implicated in self-referential processing, including the medial prefrontal cortex, precuneus, and insula. In addition, coping strategy modulated the relationship between cigarette-related brain activation and self-reported craving in a subset of these regions. These findings indicate that coping strategies that entail the generation and maintenance of self-relevant information rely upon different psychological and neurobiological mechanisms than those that are not self-focused, even when the latter incorporate information that is very similar in content. Results extend previous work examining the neural substrates of coping with craving. Given the potential mnemonic and motivational advantages associated with self-related processing, findings may have significant implications for selecting and improving techniques for helping quitting smokers resist the urge to smoke.

Keywords: coping, craving, functional MRI, self, smoking

Relapse remains the unfortunate outcome for the vast majority of individuals who attempt to quit smoking cigarettes (Piasecki, 2006). Smokers are particularly vulnerable to relapse when they encounter cigarette-related stimuli (Ferguson & Shiffman, 2009). Smoking cues are thought to be problematic largely because they elicit a strong urge or craving to smoke (Sayette & Griffin, 2010). Consistent with this view, several studies have established a link between cue-elicited craving and relapse (Ferguson & Shiffman, 2009).

Research also indicates that the likelihood of relapse during high-risk situations (e.g., when craving is elevated) can be reduced significantly through the use of strategies for coping with craving (Ferguson & Shiffman, 2009). Among the coping strategies that are used, cognitively oriented techniques (e.g., thinking about the benefits of quitting) are often employed successfully by quitting smokers when urges arise (e.g., Bliss, Garvey, & Ward, 1999). Cognitive coping methods are not fail-safe, however, as quitting smokers often succumb to temptation during high-risk situations despite reporting their use (e.g., Shiffman, Paty, Gnys, Kassel, & Hickcox, 1996). Presently, little is known about why cognitive coping strategies are effective in some situations but ineffective in others, primarily because the mechanisms through which they operate remain unclear.

In an important step toward addressing this knowledge gap, researchers have begun to identify the neurocognitive processes involved in the regulation of craving (Brody et al., 2007; Hartwell et al., 2011; Kober et al., 2010; Volkow et al., 2010). These studies indicate that the modulation of craving involves many of the same brain regions that have been implicated in the regulation of affective states more generally (Davidson, Fox, & Kalin, 2007; Ochsner & Gross, 2005). In particular, the inhibition of cue-elicited responses has been associated with increases in the activation of brain areas supporting domain-general control processes and decreases in the activation of areas supporting more circumscribed

This article was published Online First January 30, 2012.

Stephen J. Wilson, Department of Psychology, The Pennsylvania State University; Michael A. Sayette, Department of Psychology, University of Pittsburgh; Julie A. Fiez, Department of Psychology, University of Pittsburgh, and Center for the Neural Basis of Cognition.

This research was supported by Grant R01 DA02463 from the National Institute on Drug Abuse. This paper is based upon a dissertation completed by S. J. Wilson. Portions of these data were presented at the annual meeting of the Society for Neuroscience (Chicago, Illinois, November 2009). We thank Michael Pogue-Geile, Ahmad Hariri, and Richard Jennings for helpful discussion. We also are indebted to Deidra Rendenell for assistance with data collection.

Correspondence concerning this article should be addressed to Stephen J. Wilson, 438 Moore Building, The Pennsylvania State University, University Park, PA 16802. E-mail: sjw42@psu.edu

affective processes (Brody et al., 2007; Kober et al., 2010; Volkow et al., 2010).

We sought to further elucidate the mechanisms underlying coping by using functional MRI (fMRI) to examine the neural responses associated with different forms of cognitively oriented coping. Specifically, we compared a strategy that involved focusing upon the personal benefits of quitting (e.g., “I will breathe easier if I stop smoking”) to an approach that entailed focusing on information that was very similar in content but that was directed at another individual (e.g., “My son will breathe easier if I stop smoking”). These strategies were chosen for three reasons. First, similar techniques are employed spontaneously by quitting smokers (O’Connell, et al., 1998) and are taught in formal coping skills interventions (Monti & Rohsenow, 1999). Second, there is reason to predict that different mechanisms may be involved in coping that incorporates self-referential information relative to coping that is not self-focused. Specifically, a network of brain regions (e.g., medial prefrontal cortex [PFC]) appears to be involved preferentially in the processing of self-related information (Northoff, Qin, & Feinberg, 2011), perhaps even relative to information about close others (Heatherton et al., 2006). Finally, the different mechanisms supporting self-focused and other-focused coping strategies might influence their effectiveness. For instance, extensive research indicates that information is better remembered and more robust to distracters if it is related to the self than if it is processed in other ways, including by relating it to intimate others (e.g., associating it with a close friend; Symons & Johnson, 1997). Thus, determining whether self-referential versus non-self-referential coping strategies rely upon distinct processes may have significant clinical implications.

Smokers who were motivated to quit were instructed to engage in self-focused or other-focused coping during cue exposure and were given the option of smoking during the study. This manipulation was designed to create conflict between the intention to abstain and the desire to smoke. To our knowledge, this is one of the first studies to examine cue reactivity in smokers with conflicting motivations for and against smoking *and* a concomitant opportunity to smoke (see also Wilson, Sayette, & Fiez, in press). We anticipated that this tension might lead to a complex pattern of responses (Wilson, Sayette, & Fiez, 2004). In particular, we expected that self-reported urge might be influenced by coping, which presumably would attenuate craving, if successful. Importantly this urge appraisal-coping sequence is likely to occur very rapidly, rendering it nearly impossible to capture adequately, in real time, using a self-report instrument (Lazarus & Folkman, 1984). A feature of the present study is its inclusion of fMRI, which can measure these fleeting processes as they unfold in real time. That is, we believe that fMRI is especially well-suited to uncovering differences between coping strategies that may be difficult to detect via self-report.

Method

Participants

Male smokers ($N = 60$) were recruited through media advertisements. Usable data were collected from 57 participants (three were excluded due to excessive movement during scanning). Participants were required to be right-handed native English speakers

between the ages of 18 and 45, and to have smoked at least 15 cigarettes per day for 24 months. Additionally, they had to report that they were planning on quitting smoking within 2 weeks, were interested in smoking cessation treatment, and were willing to initiate a quit attempt during the experiment. Participants were randomly assigned to two coping strategy conditions (*self-focused* and *other-focused*; described in the Procedures section). Groups did not differ in age ($M = 33.6$ years, $SD = 8.5$), ethnicity (59% Caucasian, 39% African American, 2% other), cigarettes per day ($M = 20.9$, $SD = 6.0$), or nicotine dependence, as assessed using the Fagerström Test for Nicotine Dependence ($M = 4.9$, $SD = 1.6$; Heatherton, Kozlowski, Frecker, & Fagerström, 1991).

Design and Overview

Participants completed two sessions during the study, which are described in detail in the Procedures section. Those deemed eligible, based upon telephone screening, were scheduled for an initial baseline session, during which questionnaires, behavioral working memory assessments, and coping strategy training were administered. Participants then were scheduled for the fMRI-based experimental session (held within 2 weeks of the baseline session), during which they performed a working memory paradigm and cue exposure/coping task while fMRI data were collected. The experimental session was scheduled to coincide with the first day of an attempt to quit smoking. Specifically, participants were instructed to initiate a cessation attempt 12 hr prior to the onset of the experimental visit.

Materials

Questionnaires. During the baseline session, participants completed questionnaires measuring the following: current and past smoking practices (Shiffman, Paty, Kassel, Gnys, & Zettler-Segal, 1994), level of nicotine dependence (Fagerström Test for Nicotine Dependence, Heatherton, et al., 1991; Nicotine Dependence Syndrome Scale, Shiffman, Waters, & Hickcox, 2004), smoking abstinence self-efficacy (Relapse Situation Efficacy Questionnaire; Gwaltney, et al., 2001), self-consciousness (revised Self-Consciousness Scale; Scheier & Carver, 1985), trait self-control (Self Control Scale; Tangney, Baumeister, & Boone, 2004), trait positive and negative affect (Positive and Negative Affect Schedule [PANAS]; Watson, Clark, & Tellegen, 1988), and tendency to respond in a socially desirable manner (Balanced Inventory of Desirable Responding Version 6; Paulhus, 1991). In order to measure the effects of nicotine withdrawal on cognitive and emotional functioning, participants completed questionnaires measuring the following at the beginning of Session 2 (i.e., after abstaining from smoking for 12 hr): current levels of positive and negative affect (state version of the PANAS; Watson, Clark, & Tellegen, 1988) and mental energy/fatigue (State Self-Control Capacity Scale; Ciarocco, Twenge, Muraven, & Tice, 2007). Questionnaire data are not a focus of the present study and are not presented herein.

Cue exposure/coping task. Participants completed a cue-exposure procedure adapted from prior research (Wilson, Sayette, Delgado, & Fiez, 2005; Wilson, et al., in press). Runs began with a 48-s resting baseline. Next, an object was placed in the participant’s left hand, and prerecorded instructions identifying the object

were delivered via intercom. Participants held and passively viewed the object for 74 s. Participants were able to see what they were holding by way of a live video feed. Participants completed three runs, during which they held a notepad (practice), roll of tape (control cue), and cigarette in the first, second, and third run, respectively. Because exposure to smoking cues can affect responses to subsequently presented items (see Sayette, Griffin, & Sayers, 2010), the order was fixed in this sequence. The first run served as practice and was excluded from analyses. Before the third run, participants were informed that they would hold a cigarette. They were instructed to begin implementing the relevant coping strategy as soon as the cigarette was placed in their hand and to continue doing so until the run concluded. Participants also were told that they would be removed from the scanner and given an opportunity to smoke immediately following the run. Upon presentation of the cigarette, a prerecorded message was delivered informing participants that they would be removed from the scanner and would be able to smoke if they chose to do so.

Urge and affect. Participants verbally rated their urge to smoke from 0 (*absolutely no urge to smoke at all*) to 100 (*strongest urge to smoke I've ever experienced*) and their affect from 0 (*I feel very bad right now*) to 10 (*I feel very good right now*). These single-item scales have proven sensitive to a range of craving-related experiences (e.g., Sayette, Martin, Hull, Wertz, & Perrott, 2003).

Working memory tasks. Participants completed behavioral working memory assessments and an fMRI-based working memory task (a verbal n-back task, adapted from Ravizza, Delgado, Chein, Becker, & Fiez, 2004) as a part of larger study examining individual differences in working memory functioning in individuals who smoke. Results will be presented in a separate manuscript.

Procedure

Baseline assessment/training session. Participants completed a baseline session, during which they provided a carbon monoxide (CO) sample and completed questionnaires. After completing questionnaires, participants underwent a behavioral coping strategy training procedure. The training protocol integrated aspects of established coping skills interventions (Monti & Rohsenow, 1999) with procedures that have been successfully implemented in basic behavioral and neuroscientific affect regulation research (Ochsner et al., 2004). As noted, participants were randomly assigned to two coping strategy conditions (*self-focused* and *other-focused*). Those in the self-focused condition were trained to generate and maintain thoughts about the positive effects that quitting smoking would have on them personally. Participants in the other-focused condition were trained to generate and maintain thoughts about the positive effects that quitting smoking would have on a specific individual with whom they were close.

During training, participants received explicit instruction and guidance regarding the performance of the strategy that they were to implement in the experimental session. First, participants read a brief description of the relevant strategy. Participants subsequently completed several practice trials, during which they were asked to attempt to implement the appropriate strategy while being presented with smoking-related pictures previously shown to elicit robust increases in craving (Mucha, Geier, & Pauli, 1999). Fol-

lowing the completion of each practice trial, participants were asked to record what they had been thinking about during the presentation of the smoking-related picture. This material was reviewed by an experimenter, who assessed their performance and helped shape their use of the coping technique, as necessary. The experimenter also instructed participants not to use other strategies when performing the practice trials. All participants were judged to have effectively learned the relevant strategy based upon a qualitative assessment made by the experimenter.

After training, participants were scheduled for the experiment. They were told to refrain from using nicotine-containing products for at least 12 hr before arrival. In order to model the early phases of cessation, the experiment was scheduled to begin 12 hr after participants had initiated a quit attempt, as previously noted.

Experimental session. Upon arrival for the experiment, participants reported the last time they smoked, and CO was measured to check compliance with deprivation instructions. Participants had to have a CO level that was at least 50% lower than their baseline, a cutoff established based upon research using similar samples and procedures (e.g., Sayette, Loewenstein, Griffin, & Black, 2008). Participants then provided pre-cue-exposure ratings of urge and affect.

Next, participants were informed that they would be given a break during the study, at which point they would have the opportunity to smoke a cigarette, and then were placed in the scanner. After collection of anatomical images, participants completed a working memory task and then the cue exposure/coping procedure. Additional urge-and-affect ratings were collected immediately following the second and third runs of the cue exposure task while participants were still holding the tape and cigarette, respectively. Subsequently, participants were removed from the scanner and were presented with the opportunity to smoke (those who chose not to smoke were permitted to take a break). After smoking or taking a break, participants completed post-task questionnaires and were given an opportunity to participate in a follow-up study. Finally, participants were debriefed and paid.

fMRI data acquisition. Scanning was conducted using a 3-Tesla head-only Siemens Allegra magnet (Siemens Corporation, New York, NY). A 40-slice oblique-axial anatomical series ($3.125 \times 3.125 \times 3.0$ mm voxels) was acquired using a standard T2-weighted pulse sequence. Additionally, a high-resolution ($1 \times 1 \times 1$ mm voxels) structural volume was collected using a magnetization-prepared rapid gradient-echo sequence. Functional images were acquired in the same plane as the 40-slice anatomical series, with coverage limited to the 38 center slices using a one-shot echo-planar imaging pulse sequence (TR = 2000 ms, TE = 25 ms, FOV = 20 cm, flip angle = 79°). Heart rate (HR) was recorded during the acquisition of fMRI data using pulse oximetry from the right middle finger.

fMRI data analysis. Preprocessing and analysis of fMRI data was conducted using utilities from the following software packages: Analysis of Functional NeuroImages (version 2.6; Cox, 1996), Automated Image Registration (version 3.08; Woods, Cherry, & Mazziotta, 1992), the Functional Magnetic Resonance Imaging of the Brain (FMRIB) Software Library (FSL, release 4.1; Smith et al., 2004), and the NeuroImaging Software Package (version 3.5; Laboratory for Clinical Cognitive Neuroscience, University of Pittsburgh, and the Neuroscience of Cognitive Control Laboratory, Princeton University). Software integration and image

format conversion was implemented using the Functional Imaging Software Widgets graphical computing environment (Fissell et al., 2003). A series of preprocessing steps were employed to correct for artifacts and to account for individual differences in anatomy prior to analyzing fMRI data. Functional images were corrected for head motion and adjusted for drift within and between runs. Participants exhibiting movement that exceeded 3 mm or 3° were excluded from subsequent analysis (three participants were excluded on this basis, as previously noted). Anatomical images from each participant were coregistered to a common reference anatomy using a 6-parameter rigid-body automated registration algorithm. The transformation matrix generated during this step was then applied to the participant's functional images. Subsequently, functional images were globally mean-normalized and smoothed using a three-dimensional (3D) Gaussian filter (4-mm full width at half maximum). Group-based statistical maps were transformed into MNI stereotaxic space (FSL's MNI 152; T1, 1 × 1 × 1 mm) for anatomical localization.

The following steps were used to analyze data from the cue exposure/coping task. First, predictors for each cue type (i.e., control and cigarette) were entered into a general linear model to obtain beta weights for each participant. As in previous work (Wilson et al., 2005), data collected during the final 48 s of the control and cigarette-cue epochs were included in analyses; data collected during the initial 26 s were excluded to allow for stabilization of responses associated with prerecorded object identification messages. Beta weights were divided by the estimated run baseline to convert them to units of percent change and were entered into a second-level 2 (coping strategy) × 2 (cue) mixed-model ANOVA. We also examined the relationship between activation during cigarette cue exposure and other cue-reactivity measures (self-reported urge, HR) using whole-brain multiple regression with three covariates: coping strategy (dummy coded), the relevant cue-reactivity measure (self-reported urge or HR; mean centered), and the coping strategy by cue-reactivity interaction (product of the coping strategy and urge/HR covariates).

Monte Carlo simulations indicated that a combined per-voxel threshold of $p < .005$ and cluster-extent threshold of 11 or more contiguous voxels would yield a corrected clusterwise false positive rate of $p < .05$. These parameters were applied to all statistical maps.

Results

Smoking Behavior and Quit Interest

Five participants in the self-focused condition and eight participants in the other-focused condition chose not to smoke during the experiment; the remaining 44 participants smoked when given the opportunity to do so. The relationship between coping strategy and choice was not significant, $\chi^2(1, N = 57) = .77, p > .5$. Participants rated their current interest in quitting at the conclusion of the experiment from 1 (*not at all interested*) to 10 (*extremely interested*). Ratings from two participants were missing; results include data from the remaining 55. Participants in the self-focused ($M = 8.42, SD = 1.27$) and other-focused ($M = 8.83, SD = 1.20$) conditions did not differ in their reported interest in quitting, $t(53) = 1.22, p > .2, \eta^2 = .03$.

Urge and Affect

Mean urge and affect levels are presented in Table 1. A 2 (coping strategy) × 3 (time) mixed-model ANOVA with the three urge ratings as a repeated variable did not yield significant effects ($ps > .2$). Similarly, an ANOVA conducted with self-reported affect as the dependent measure did not produce significant results ($ps > .2$).

Unlike most prior smoking cue-reactivity studies, participants in this study were motivated to quit smoking and were asked to engage in coping during cue exposure. We speculated that a significant proportion of participants might not have exhibited significant increases in urge because of these unique methodological features (i.e., because coping was effective or because they were reluctant to report high levels of urge; see Wertz & Sayette, 2001). As presented in Table 2 and Figure 1, we observed substantial variability in cue-elicited self-reported urge, with similar patterns exhibited by both coping conditions. In order to explore the possibility that some participants may have been reluctant to endorse elevated craving, we examined the relationship between self-reported urge during cigarette cue exposure and scores on a questionnaire assessing the tendency to respond in a socially desirable manner (Balanced Inventory of Desirable Responding Version 6 [BIDR-6]; Paulhus, 1991). Craving ratings provided at the conclusion of the cigarette cue exposure period negatively correlated with scores on the BIDR-6, $r(57) = -.37, p < .01$,

Table 1
Mean (SD) for Urge, Affect, and Heart Rate

	Full sample ($n = 57$)	Self-focused condition ($n = 28$)	Other-focused condition ($n = 29$)
Affect-baseline	7.1 (1.7)	7.1 (1.9)	7.0 (1.7)
Affect-control cue	7.0 (2.3)	6.8 (2.7)	7.2 (1.8)
Affect-cigarette cue	7.0 (2.3)	6.7 (2.7)	7.3 (1.9)
Urge-baseline	62.5 (25.7)	64.5 (22.6)	60.6 (28.6)
Urge-control cue	64.3 (28.8)	61.1 (30.2)	67.4 (27.6)
Urge-cigarette cue	64.0 (31.7)	60.6 (33.1)	67.2 (30.5)
Heart rate-control cue	60.2 (8.1)	61.0 (7.9)	59.6 (8.4)
Heart rate-cigarette cue	61.2 (8.6)	61.9 (8.4)	60.6 (8.8)

Note. Heart rate is presented in beats per minute.

Table 2
Percentage of Participants Reporting an Increase, No Change, or a Decrease in Craving During Cigarette Relative to Control Cue Exposure

	Full sample (<i>n</i> = 57)	Self-focused condition (<i>n</i> = 28)	Other-focused condition (<i>n</i> = 29)
Increase	47	50	45
No change	26	29	24
Decrease	26	21	31

$\eta^2 = .14$, suggesting that urge ratings may have at least in part reflected demand characteristics associated with the study (cf. Sayette & Parrott, 1999).

Heart Rate

Due to technical error, data were not collected from six participants (five self-focused, one other-focused). Using available data, we conducted a 2 (coping strategy) \times 2 (cue) ANOVA, with HR (beats/min) averaged across the time period during which the control and cigarette cues were held as the dependent variable. We observed a significant main effect of cue, with greater HR during the cigarette cue than the control cue, $F(1, 49) = 7.84, p < .01, \eta^2 = .14$ (see Table 1). None of the remaining effects were significant.

fMRI Results

Main effect of cue. Regions exhibiting a main effect of cue are presented in Table 3. Activation was greater during the presentation of the cigarette cue than the control cue in several areas, including the PFC, anterior and posterior cingulate, thalamus, and cerebellum. Greater activation during the control cue relative to the cigarette was observed bilaterally in the superior temporal gyrus.

Coping Strategy \times Cue interaction. A significant Coping Strategy \times Cue interaction was observed in several regions implicated in self-referential processing, including midline cortical structures (medial PFC, dorsal and rostral anterior cingulate [ACC], and precuneus) and bilaterally in the insula (see Table 4 and Figure 2). The large medial frontal region that exhibited a Coping Strategy \times Cue interaction partially overlapped with a midline region that showed a main effect of cue. (Specifically, 12% of the voxels in the medial PFC/dorsal ACC demonstrating a main effect also displayed an interaction.) The effect of cue was examined separately for the self-focused and other-focused conditions to characterize the interaction for each region. As presented in Table 4, activation during the cigarette cue was greater than that during the control cue for participants in the self-focused condition in each region. In contrast, activation during the control cue either did not differ from, or was significantly greater than, activation during the cigarette cue for participants in the other-focused condition in these areas.

Modulation of activation/cue-reactivity correlations by coping strategy. A significant relationship between cigarette-related activation and urge ratings during cigarette cue exposure was observed in the right superior frontal gyrus (MNI coordinates: $x = 26, y = -2, z = 62$; Brodmann Area [BA] 6; size = 322

mm³; average F-ratio = 11.76) and right dorsolateral PFC ($x = 52, y = 33, z = 14$; BA 46; size = 322 mm³; average F-ratio = 10.89). Considering the sample as a whole, greater activation was associated with lower self-reported urge to smoke during cigarette cue exposure for both regions of interest (ROIs). For each region, however, this relationship was driven by a robust negative association between cue-elicited urge and brain activation for the other-focused group (right superior frontal gyrus: $r[29] = -.64, p < .001, \eta^2 = .41$; right dorsolateral PFC: $r[29] = -.63, p < .001, \eta^2 = .40$). In contrast, for the self-focused group, cue-elicited urge was positively related to activation of the dorsolateral PFC ($r[28] = .40, p < .04, \eta^2 = .16$) and unrelated to activation of the superior frontal gyrus ($r[28] = .27, p = .16, \eta^2 = .07$).

Of particular interest, a significant relationship between cue-elicited activation and the interaction between coping strategy and smoking urge was observed in the left inferior parietal lobule, bilateral insula, and right inferior frontal gyrus (see Table 5 and Figure 2). As shown in Figure 2, these regions fall close to, or overlap with, those exhibiting a significant coping strategy by cue interaction (the right insula/inferior frontal gyrus and left insula ROIs identified in the multiple regression analysis overlap with 66% and 16% of the voxels exhibiting a significant coping strategy by cue interaction, respectively). Cue-elicited activation of each region was positively correlated with urge ratings during cue exposure for the self-focused group. In contrast, activation was unrelated to (right insula) or negatively correlated with (inferior parietal lobule, left insula, right inferior frontal gyrus/insula) urge for the other-focused group. (As indicated in Table 5, the effect was marginally significant [$p < .1$] for the left insula and right inferior frontal gyrus/insula.)

We observed a negative correlation between cigarette-related activation and HR during cigarette cue exposure in the right anteroventral PFC for both coping groups ($x = 37, y = 46, z = -10$; BA 10/47; size = 557 mm³; average F-ratio = 15.68). We also observed a significant relationship between cue-elicited acti-




Figure 1. Scatterplot of self-reported urge during cigarette cue exposure for the self-focused and other-focused groups.

Table 3
Brain Regions Exhibiting a Significant Main Effect of Cue

Region	BA	Size (mm ³)	MNI coordinates			Average <i>F</i> ratio
			x	y	z	
Cigarette > Control						
ACC/superior frontal g/SMA	6/32	2168	-4	7	55	12.41
L middle frontal g (dorsolateral PFC)	9	322	-43	3	41	10.76
Posterior cingulate g	29	1260	8	-28	27	9.94
L middle frontal g	10	938	-34	45	25	11.39
R parahippocampal g	34	352	18	-24	-20	12.55
L caudate nucleus/thalamus		1846	-18	-2	20	11.35
R thalamus		586	7	-19	15	10.36
Cerebellum		410	33	-62	-30	12.47
Cerebellum		469	-30	-58	-33	10.49
Control > Cigarette						
L superior temporal g	22	3105	-60	-26	6	14.54
R superior temporal g	22	1289	60	-13	0	13.08

Note. Stereotaxic coordinates are given for local maxima of activation cluster in MNI atlas space. ACC = anterior cingulate cortex; BA = Brodmann's area; g = gyrus; L = left hemisphere; PFC = prefrontal cortex; R = right hemisphere; SMA = supplementary motor area.

vation and the interaction between coping strategy and HR in the middle and inferior occipital gyri bilaterally (left hemisphere: $x = -26$, $y = -82$, $z = -9$; BA 18; size = 674 mm³; average *F*-ratio = 10.63; right hemisphere: $x = 30$, $y = -87$, $z = -1$; BA 18; size = 674 mm³; average *F*-ratio = 11.22). Cue-elicited activation of both regions was negatively correlated with HR during cigarette exposure for participants in the self-focused condition, while cigarette-related activation and HR were not correlated for participants in the other-focused condition.

Discussion

We examined neural activity in quitting-motivated male cigarette smokers engaging in self-focused or other-focused coping while being presented with a cigarette cue and an opportunity to smoke. Several brain regions exhibited significant main effects due to cue exposure. Consistent with previous research, cigarette-related increases in activation were observed for both groups in the middle frontal gyrus (e.g., Due, Huettel, Hall, & Rubin, 2002; McBride, Barrett, Kelly, Aw, & Dagher, 2006), ACC (e.g., Brody

et al., 2007; Wagner, Dal Cin, Sargent, Kelley, & Heatherton, 2011), posterior cingulate (e.g., McClernon, Kozink, & Rose, 2008; Wilson et al., 2005), dorsal striatum (e.g., McClernon, Kozink, Lutz, & Rose, 2009; Yalachkov, Kaiser, & Naumer, 2009), parahippocampal gyrus (e.g., Janes et al., 2010; Smolka et al., 2006), and cerebellum (e.g., McClernon et al., 2008; Wilson, et al., in press).

Of primary interest, we also found that coping strategy affected cue-related activation in several brain areas. Quitting smokers who were told to focus on the personal benefits of smoking cessation, but not those told to focus on the benefits that cessation would have on someone close to them, exhibited significant cue-related activation of the medial PFC, precuneus, and dorsal and rostral ACC (the dorsal ACC region partially overlapping with an area exhibiting a main effect of cue). These findings are consistent with extensive evidence implicating these midline cortical structures as key brain areas supporting self-referential processing (Cavanna & Trimble, 2006; Heatherton et al., 2006; Northoff et al., 2006). Also of note, only the self-focused coping condition was associated with

Table 4
Brain Regions Exhibiting a Significant Coping Strategy \times Cue Interaction

Region	BA	Size (mm ³)	MNI coordinates			Average <i>F</i> ratio	Self-focused condition ($n = 28$)	Other-focused condition ($n = 29$)
			x	y	z			
L precuneus	7	439	-12	-69	46	11.17	Cigarette > Control**	Control > Cigarette*
Medial frontal g/dorsal ACC	8/32	1230	-1	32	45	11.70	Cigarette > Control**	Control > Cigarette*
L inferior parietal lobule	40	410	-48	-45	36	9.67	Cigarette > Control**	ns
L middle frontal g (rostral PFC)	10	615	-37	38	26	10.65	Cigarette > Control**	ns
Rostral ACC	24	381	-1	30	6	10.21	Cigarette > Control**	Control > Cigarette*
R insula/inferior frontal g	13/45/47	527	34	24	6	11.23	Cigarette > Control**	ns
L insula/inferior frontal g/superior temporal g	13/47/22	2490	-36	18	0	11.20	Cigarette > Control**	Control > Cigarette**
Cerebellum		352	-12	-77	-28	9.63	Cigarette > Control**	Control > Cigarette*

Note. Stereotaxic coordinates are given for local maxima of activation cluster in MNI atlas space. ACC = anterior cingulate cortex; BA = Brodmann's area; g = gyrus; L = left hemisphere; ns = no significant effect of cue; PFC = prefrontal cortex; R = right hemisphere.

* $p < .05$. ** $p < .01$.




Figure 2. Row 1: Regions exhibiting a significant coping strategy (self-focused vs. other-focused) by cue (control, cigarette) interaction. Row 2: Regions exhibiting a significant relationship between cigarette-related activation and the interaction of coping strategy and urge to smoke during cigarette cue exposure. The numbers below each column denote the distance (mm) of the brain slice from the anterior commissure–posterior plane in MNI stereotaxic space. Brain slices are right–left reversed. ACC = anterior cingulate cortex; IFG = inferior frontal gyrus; INS = insula; IPL = inferior parietal lobule; MFG = medial frontal gyrus; Pcu = precuneus.

cue-related activation of the anterior insula, a region that has received increasing attention from addiction researchers (Garavan, 2010; Naqvi & Bechara, 2010; Naqvi, Rudrauf, Damasio, & Bechara, 2007) and that appears to play an important role in self-related processing (Craig, 2009; Feinberg, 2011; Northoff et al., 2011). Presumably, the greater activation of this set of regions (particularly medial cortical structures) by the self-focused, relative to the other-focused, group reflects the higher degree of self-related processing (e.g., self-focused attention) utilized by the former during cue exposure. Such effects may have important clinical implications, as recent findings suggest that the success of smoking cessation interventions relates to the degree to which they engender a sense of personal relevance (Chua et al., 2011; Chua, Liberzon, Welsh, & Strecher, 2009).

We found additional evidence of differences between the coping strategies upon examining the relationship between brain activation and other cue-reactivity measures. For the other-focused group, smoking urge was negatively associated with cue-related

activation of the dorsolateral PFC and superior frontal gyrus, regions that recently have been linked to the modulation of craving (Hartwell et al., 2011; Kober et al., 2010; Rose et al., 2011). In contrast, self-reported urge was positively associated with cue-related activation of the dorsolateral PFC, but unrelated to activation of the superior frontal gyrus, for the self-focused group. Divergent associations between cue-elicited urge and brain activation as a function of coping strategy were observed in several additional regions. Specifically, urge was positively related to cue-elicited activation of the left inferior parietal lobule, bilateral insula, and right inferior frontal gyrus for the self-focused group, but negatively correlated with or unrelated to activation of these areas for the other-focused group. Notably, these areas overlap with those that demonstrated a significant coping strategy by cue interaction, suggesting that the different neural activation patterns exhibited by the self-focused and other-focused groups are behaviorally relevant. Effects observed in the insula, which appears to play a critical role in drug craving (Garavan, 2010; Naqvi &

Table 5

Coping Strategy Moderates the Relationship Between Brain Activation and Self-Reported Urge During Cigarette Cue Exposure

Region	BA	Size (mm ³)	MNI coordinates			Average <i>F</i> ratio	Self-Focused Condition (<i>n</i> = 28)	Other-Focused Condition (<i>n</i> = 29)
			x	y	z			
L inferior parietal lobule	40	615	−59	−36	33	10.89	Positive ^{***}	Negative ^{***}
L insula	13	352	−43	7	2	10.11	Positive ^{**}	Negative [*]
R insula	13	527	45	10	1	10.30	Positive ^{***}	ns
R inferior frontal g/insula	47/13	439	40	24	1	10.56	Positive ^{**}	Negative [*]

Note. Stereotaxic coordinates are given for local maxima of activation cluster in Montreal Neurological Institute (MNI) atlas space. BA = Brodmann's area; g = gyrus; L = left hemisphere; Negative = negative correlation between brain activation and self-reported urge during cigarette cue exposure; ns = no significant relationship between brain activation and self-reported urge during cigarette cue exposure; Positive = positive correlation between brain activation and self-reported urge during cigarette cue exposure; R = right hemisphere.

* $p < .10$. ** $p < .01$. *** $p < .001$.

Bechara, 2010; Naqvi et al., 2007), are of particular interest. Further research is needed to determine whether the observed effects are associated with strategy-related differences in the regulation of the interoceptive processes mediated by the insula, differences in the use of such processes in support of coping, or some other mechanism.

Additionally, HR was differentially coupled with the activation of bilateral extrastriate visual cortical regions as a function of coping condition (i.e., cue-elicited activation in the visual cortex was negatively correlated with HR during cue exposure for the self-focused group, while cigarette-related activation and HR were not related for the other-focused group). This result may have clinical significance, as Brody and colleagues (2007) found that treatment-seeking smokers exhibited decreased cue-elicited activation of the visual cortex when they were told to resist their craving compared with when they were told to allow themselves to crave. Additional research is needed to elucidate the functional significance of this pattern, as cue-elicited changes in HR can reflect a variety of influences (see Carter & Tiffany, 1999; Sayette et al., 2000).

Taken together, our findings suggest that coping involving the generation and maintenance of self-relevant information rely upon different psychological and neurobiological mechanisms than those that are not self-focused, even when the latter incorporate information that is very similar in content. As noted, the different processes underpinning self-focused and non-self-focused cognitive coping techniques may have implications for their relative efficacy. Information is better remembered and more robust to distracters if it is related to the self than if it is processed in other ways (Symons & Johnson, 1997). To the extent that such effects extend to the domain of coping, attempting to focus on their personal reasons for quitting may provide a mnemonic advantage relative to trying to think of other potential sources of motivation for those attempting to discontinue smoking. Interestingly, recent studies have found that some of the same regions identified in the current investigation (medial PFC and precuneus/posterior cingulate) were more strongly activated by smoking cessation messages that contained personalized material than those containing generic information (Chua et al., 2009), and that the degree to which such regions were engaged by personally tailored messages predicted the odds of quitting smoking 4 months later (Chua et al., 2011). These findings highlight personal relevance (which may influence motivation to change behavior) as an important factor in determining the success of smoking cessation interventions, offering another potential benefit of self-focused relative to non-self-focused coping strategies (i.e., in addition to the potential mnemonic advantages afforded by the former).

It is important to note, however, that although we have focused on the potential advantages associated with coping that heavily involves self-related processing, we did not find direct evidence that the self-focused and other-focused strategies differed in effectiveness, leaving open the possibility that the strategies were equally efficacious (or even that the other-focused strategy was superior to the self-focused strategy). As noted, the majority of participants in each condition chose to smoke when given the opportunity to do so. In addition, both groups failed to exhibit significant increases in self-reported urge during cue exposure. The lack of self-reported urge effects may relate to key differences between the methods employed in the current investigation and the

procedures typically used in prior research. Specifically, participants were asked to engage in coping while being exposed to a cigarette cue in the present study, whereas participants in prior studies generally were not asked to engage in any form of self-regulation. It is possible that the lack of urge increases during cue exposure in the present study may reflect the influence of coping (i.e., which attenuated increases in urge for some). As noted, this process is likely to occur rapidly (Lazarus & Folkman, 1984), highlighting the utility of fMRI for revealing strategy-related differences that were not detected through the assessment of self-report. The lack of urge effects may also relate, in part, to the influence of experimental demand (cf. Sayette & Parrott, 1999). Still another possibility is that participants may not have reported increases in urge because they were motivated to quit smoking (see Wertz & Sayette, 2001). Regardless, the significant variability in urge exhibited by participants allowed us to examine relationships between cue-elicited brain activity and craving.

Recent data from our laboratory indicate that quitting smokers do endorse cue-elicited increases in urge, at least under certain conditions. Specifically, in a recent fMRI study utilizing the same cue exposure procedure, quitting-motivated smokers who were told that they could smoke—and were not asked to engage in coping—reported a mean urge level of 77 on a 0–100 scale while holding a cigarette, which was significantly greater than the mean urge of 68 that they reported while holding a control cue (Wilson, et al., in press). These results support the idea that coping influenced urge reporting in the current study.

Additional limitations should be mentioned. This study included only male smokers. This decision was based upon research demonstrating that male and females exhibit different patterns of neural activation during the affect regulation (Domes et al., 2010; McRae, Ochsner, Mauss, Gabrieli, & Gross, 2008) and drug cue exposure (Kilts, Gross, Ely, & Drexler, 2004; McClernon et al., 2008). We sought to maximize our ability to detect effects of interest by excluding potential sex differences. Additional research is needed to determine whether or not findings generalize to female smokers.

It also is worth noting that, while cue-related activation was observed in several brain areas identified in previous research, we failed to find cue-elicited activation in other regions thought to play an important role in drug addiction, including the amygdala and ventral striatum (Koob & Volkow, 2010). Activation in one or both of these brain areas has been observed in several prior studies (e.g., David et al., 2005; Franklin et al., 2009; Franklin, Wang, Li, et al., 2011; Franklin, Wang, Suh, et al., 2011; Franklin et al., 2007; Stippekohl et al., 2010). It is possible that the absence of effects in these regions relate to the use of coping by participants in the present study (e.g., see Volkow et al., 2010). However, given that we failed to observe main effects in these regions in prior studies that did not involve coping (Wilson et al., 2005; Wilson, et al., in press), it is likely that the discrepancy between our results and prior findings is at least in part due to factors other than coping, such as differences in imaging technique (e.g., blood-oxygen-level-dependent vs. arterial spin labeling perfusion fMRI) or cue exposure methods (e.g., the use of a single, relatively static smoking cue vs. the use of more dynamic or multiple cues). Future research is needed to explore these possibilities.

Notwithstanding these potential limitations, the present findings highlight the importance of examining the neurocognitive mech-

anisms supporting different cognitively oriented coping strategies. In addition to advancing understanding of the coping process, investigating how various coping techniques operate may facilitate efforts to select and improve techniques for helping quitting smokers resist temptation. To our knowledge, this is the first study to demonstrate that self-referential and non-self-referential coping strategies are associated with different patterns of brain activation. By helping to establish key linkages between coping with craving, the emerging neuroscience literature on the neurobiological underpinnings of self-referential processing (Northoff et al., 2011), and the extensive cognitive psychology literature on self-referential encoding (Symons & Johnson, 1997), the present findings also point toward useful targets for future research. Given the potential mnemonic and motivational advantages associated with self-related processing, additional research utilizing more extensive training procedures to explore the clinical concomitants of self-referential versus non-self-referential strategies for regulating urge may be particularly fruitful. In addition, research exploring factors that may serve to differentially moderate the effectiveness of self-focused relative to non-self-focused coping techniques, such as genetically driven variation in the functioning of regions more strongly linked to one approach to coping than the other (e.g., see Franklin et al., 2009; Franklin, Wang, Li, et al., 2011), would be useful.

Results from the current study also have important methodological implications. As noted above, naturalistic studies suggest that quitting smokers often spontaneously implement self-referential and non-self-referential coping strategies (O'Connell, et al., 1998). To the extent that the use of such techniques differs across individuals, or across time within individuals, it will be important for researchers to account for coping strategy as a potential source of variance (cf. Hartwell et al., 2011). More generally, our findings suggests that it is feasible to study cue reactivity and coping in smokers under conditions that have not previously been investigated in detail. Specifically, an attempt was made to create a high degree of conflict between the intention to abstain and the urge to smoke by selecting participants who were motivated to quit smoking and subsequently presenting them with smoking cues and an opportunity to smoke. The current study provides preliminary support for the idea that this unique motivational state can be produced under controlled laboratory conditions. As the results from such research stand to greatly inform our understanding of the ambivalence associated with addiction and relapse, additional work using this approach is indicated. Finally, this research highlights the utility of brain imaging approaches to capture rapidly emerging processing associated with urge appraisal and coping appraisal that are difficult to assess unobtrusively and in real time using self-report instruments.

References

- Bliss, R. E., Garvey, A. J., & Ward, K. D. (1999). Resisting temptations to smoke: Results from within-subjects analyses. *Psychology of Addictive Behaviors, 13*, 143–151. doi:10.1037/0893-164X.13.2.143
- Brody, A. L., Mandelkern, M. A., Olmstead, R. E., Jou, J., Tiongson, E., Allen, V., . . . Cohen, M. S. (2007). Neural substrates of resisting craving during cigarette cue exposure. *Biological Psychiatry, 62*, 642–651. doi:10.1016/j.biopsych.2006.10.026
- Carter, B. L., & Tiffany, S. T. (1999). Meta-analysis of cue-reactivity in addiction research. *Addiction, 94*, 327–340. doi:10.1046/j.1360-0443.1999.9433273.x
- Cavanna, A. E., & Trimble, M. R. (2006). The precuneus: A review of its functional anatomy and behavioural correlates. *Brain: A Journal of Neurology, 129*, 564–583. doi:10.1093/brain/awl004
- Chua, H. F., Ho, S. S., Jasinska, A. J., Polk, T. A., Welsh, R. C., Liberzon, I., & Strecher, V. J. (2011). Self-related neural response to tailored smoking-cessation messages predicts quitting. *Nature Neuroscience, 14*, 426–427. doi:10.1038/nn.2761
- Chua, H. F., Liberzon, I., Welsh, R. C., & Strecher, V. J. (2009). Neural correlates of message tailoring and self-relatedness in smoking cessation programming. *Biological Psychiatry, 65*, 165–168. doi:10.1016/j.biopsych.2008.08.030
- Ciarocco, N. J., Twenge, J. M., Muraven, M., & Tice, D. M. (2007). *The state self-control capacity scale: Reliability, validity, and correlations with physical and psychological stress*. Unpublished manuscript.
- Cox, R. W. (1996). AFNI: Software for analysis and visualization of functional resonance neuroimages. *Computational and Biomedical Research, 29*, 162–173. doi:10.1006/cbmr.1996.0014
- Craig, A. D. (2009). How do you feel—now? The anterior insula and human awareness. *Nature Reviews Neuroscience, 10*, 59–70. doi:10.1038/nrn2555
- David, S. P., Munafo, M. R., Johansen-Berg, H., Smith, S. M., Rogers, R. D., Matthews, P. M., & Walton, R. T. (2005). Ventral striatum/nucleus accumbens activation to smoking-related pictorial cues in smokers and nonsmokers: A functional magnetic resonance imaging study. *Biological Psychiatry, 58*, 488–494. doi:10.1016/j.biopsych.2005.04.028
- Davidson, R. J., Fox, A., & Kalin, N. H. (2007). Neural bases of emotion regulation in nonhuman primates and humans. In J. J. Gross (Ed.), *Handbook of emotion regulation* (pp. 47–68). New York, NY: Guilford Press.
- Domes, G., Schulze, L., Bottger, M., Grossmann, A., Hauenstein, K., Wirtz, P. H., . . . Herpertz, S. C. (2010). The neural correlates of sex differences in emotional reactivity and emotion regulation. *Human Brain Mapping, 31*, 758–769. doi:10.1002/hbm.20903
- Due, D. L., Huettel, S. A., Hall, W. G., & Rubin, D. C. (2002). Activation in mesolimbic and visuospatial neural circuits elicited by smoking cues: Evidence from functional magnetic resonance imaging. *The American Journal of Psychiatry, 159*, 954–960. doi:10.1176/appi.ajp.159.6.954
- Feinberg, T. E. (2011). The nested neural hierarchy and the self. *Consciousness and Cognition: An International Journal, 20*, 4–15. doi:10.1016/j.concog.2010.09.016
- Ferguson, S. G., & Shiffman, S. (2009). The relevance and treatment of cue-induced cravings in tobacco dependence. *Journal of Substance Abuse Treatment, 36*, 235–243. doi:10.1016/j.jsat.2008.06.005
- Fissell, K., Tseytlin, E., Cunningham, D., Iyer, K., Carter, C. S., Schneider, W., & Cohen, J. D. (2003). A graphical computing environment for neuroimaging analysis. *Neuroinformatics, 1*, 111–126. doi:10.1385/NI:1:1:111
- Franklin, T. R., Lohoff, F. W., Wang, Z., Sciortino, N., Harper, D., Li, Y., . . . Childress, A. R. (2009). DAT genotype modulates brain and behavioral responses elicited by cigarette cues. *Neuropsychopharmacology, 34*, 717–728. doi:10.1038/npp.2008.124
- Franklin, T. R., Wang, Z., Li, Y., Suh, J. J., Goldman, M., Lohoff, F. W., . . . Childress, A. R. (2011). Dopamine transporter genotype modulation of neural responses to smoking cues: Confirmation in a new cohort. *Addiction Biology, 16*, 308–322. doi:10.1111/j.1369-1600.2010.00277.x
- Franklin, T. R., Wang, Z., Suh, J. J., Hazan, R., Cruz, J., Li, Y., . . . Childress, A. R. (2011). Effects of varenicline on smoking cue-triggered neural and craving responses. *Archives of General Psychiatry, 68*, 516–526. doi:10.1001/archgenpsychiatry.2010.190
- Franklin, T. R., Wang, Z., Wang, J., Sciortino, N., Harper, D., Li, Y., . . . Childress, A. R. (2007). Limbic activation to cigarette smoking cues

- independent of nicotine withdrawal: A perfusion fMRI study. *Neuropsychopharmacology*, 32, 2301–2309. doi:10.1038/sj.npp.1301371
- Garavan, H. (2010). Insula and drug cravings. *Brain Structure & Function*, 214, 593–601. doi:10.1007/s00429-010-0259-8
- Gwaltney, C. J., Shiffman, S., Norman, G. J., Paty, J. A., Kassel, J. D., Gnys, M., . . . Balabanis, M. (2001). Does smoking abstinence self-efficacy vary across situations? Identifying context-specificity within the Relapse Situation Efficacy Questionnaire. *Journal of Consulting and Clinical Psychology*, 69, 516–527. doi:10.1037/0022-006X.69.3.516
- Hartwell, K. J., Johnson, K. A., Li, X., Myrick, H., Lematty, T., George, M. S., & Brady, K. T. (2011). Neural correlates of craving and resisting craving for tobacco in nicotine dependent smokers. *Addiction Biology*, 16, 654–666.
- Heatherington, T. F., Kozlowski, L. T., Frecker, R. C., & Fagerström, K. O. (1991). The Fagerstrom Test for Nicotine Dependence: A revision of the Fagerström Tolerance Questionnaire. *British Journal of Addiction*, 86, 1119–1127. doi:10.1111/j.1360-0443.1991.tb01879.x
- Heatherington, T. F., Wyland, C. L., Macrae, C. N., Demos, K. E., Denny, B. T., & Kelley, W. M. (2006). Medial prefrontal activity differentiates self from close others. *Social Cognition and Affective Neuroscience*, 1, 18–25. doi:10.1093/scan/ns1001
- Janes, A. C., Pizzagalli, D. A., Richardt, S., Frederick Bde, B., Holmes, A. J., Sousa, J., . . . Kaufman, M. J. (2010). Neural substrates of attentional bias for smoking-related cues: An fMRI study. *Neuropsychopharmacology*, 35, 2339–2345. doi:10.1038/npp.2010.103
- Kilts, C. D., Gross, R. E., Ely, T. D., & Drexler, K. P. G. (2004). The neural correlates of cue-induced craving in cocaine-dependent women. *The American Journal of Psychiatry*, 161, 233–241. doi:10.1176/appi.ajp.161.2.233
- Kober, H., Mende-Siedlecki, P., Kross, E. F., Weber, J., Mischel, W., Hart, C. L., & Ochsner, K. N. (2010). Prefrontal-striatal pathway underlies cognitive regulation of craving. *Proceedings of the National Academy of Sciences of the United States of America*, 107, 14811–14816. doi:10.1073/pnas.1007779107
- Koob, G. F., & Volkow, N. D. (2010). Neurocircuitry of addiction. *Neuropsychopharmacology*, 35, 217–238. doi:10.1038/npp.2009.110
- Lazarus, R. S., & Folkman, S. (1984). *Stress, appraisal, and coping*. New York, NY: Springer.
- McBride, D., Barrett, S. P., Kelly, J. T., Aw, A., & Dagher, A. (2006). Effects of expectancy and abstinence on the neural response to smoking cues in cigarette smokers: An fMRI study. *Neuropsychopharmacology*, 31, 2728–2738. doi:10.1038/sj.npp.1301075
- McClernon, F. J., Kozink, R. V., Lutz, A. M., & Rose, J. E. (2009). 24-h smoking abstinence potentiates fMRI-BOLD activation to smoking cues in cerebral cortex and dorsal striatum. *Psychopharmacology*, 204, 25–35. doi:10.1007/s00213-008-1436-9
- McClernon, F. J., Kozink, R. V., & Rose, J. E. (2008). Individual differences in nicotine dependence, withdrawal symptoms, and sex predict transient fMRI-BOLD responses to smoking cues. *Neuropsychopharmacology*, 33, 2148–2157. doi:10.1038/sj.npp.1301618
- McRae, K., Ochsner, K. N., Mauss, I. B., Gabrieli, J. D., & Gross, J. J. (2008). Gender differences in emotion regulation: An fMRI study of cognitive reappraisal. *Group Processes & Intergroup Relations*, 11, 143–162. doi:10.1177/1368430207088035
- Monti, P. M., & Rohsenow, D. J. (1999). Coping-skills training and cue-exposure therapy in the treatment of alcoholism. *Alcohol Research & Health*, 23, 107–115.
- Mucha, R. F., Geier, A., & Pauli, P. (1999). Modulation of craving by cues having differential overlap with pharmacological effect: Evidence for cue approach in smokers and social drinkers. *Psychopharmacology*, 147, 306–313. doi:10.1007/s002130051172
- Naqvi, N. H., & Bechara, A. (2010). The insula and drug addiction: An interoceptive view of pleasure, urges, and decision-making. *Brain Structure & Function*, 214, 435–450. doi:10.1007/s00429-010-0268-7
- Naqvi, N. H., Rudrauf, D., Damasio, H., & Bechara, A. (2007). Damage to the insula disrupts addiction to cigarette smoking. *Science*, 315, 531–534. doi:10.1126/science.1135926
- Northoff, G., Heinzel, A., de Greck, M., Bermpohl, F., Dobrowolny, H., & Panksepp, J. (2006). Self-referential processing in our brain—A meta-analysis of imaging studies on the self. *Neuroimage*, 31, 440–457.
- Northoff, G., Qin, P., & Feinberg, T. E. (2011). Brain imaging of the self—Conceptual, anatomical and methodological issues. *Consciousness and Cognition*, 20, 52–63.
- Ochsner, K. N., & Gross, J. J. (2005). The cognitive control of emotion. *Trends in Cognitive Sciences*, 9, 242–249. doi:10.1016/j.tics.2005.03.010
- Ochsner, K. N., Ray, R. D., Cooper, J. C., Robertson, E. R., Chopra, S., Gabrieli, J. D. E., & Gross, J. J. (2004). For better or for worse: Neural systems supporting the cognitive down- and up-regulation of negative emotion. *Neuroimage*, 23, 483–499. doi:10.1016/j.neuroimage.2004.06.030
- O’Connell, K. A., Gerkovich, M. M., Cook, M. R., Shiffman, S., Hickcox, M., & Kakolewski, K. E. (1998). Coping in real time: Using ecological momentary assessment techniques to assess coping with the urge to smoke. *Research in Nursing & Health*, 21, 487–497. doi:10.1002/(SICI)1098-240X(199812)21:6<487::AID-NUR3>3.0.CO;2-G
- Paulhus, D. L. (1991). Measurement and control of response bias. In J. P. Robinson, P. R. Shaver, & L. S. Wrightsman (Eds.), *Measures of personality and social psychological attitudes* (pp. 17–59). New York, NY: Academic Press.
- Piasecki, T. M. (2006). Relapse to smoking. *Clinical Psychology Review*, 26, 196–215. doi:10.1016/j.cpr.2005.11.007
- Ravizza, S. M., Delgado, M. R., Chein, J. M., Becker, J. T., & Fiez, J. A. (2004). Functional dissociations within the inferior parietal cortex in verbal working memory. *Neuroimage*, 22, 562–573. doi:10.1016/j.neuroimage.2004.01.039
- Rose, J. E., McClernon, F. J., Froeliger, B., Behm, F. M., Preud’homme, X., & Krystal, A. D. (2011). Repetitive transcranial magnetic stimulation of the superior frontal gyrus modulates craving for cigarettes. *Biological Psychiatry*, 70, 794–799.
- Sayette, M. A., & Griffin, K. M. (2011). Self-regulatory failure and addiction. In K. D. Vohs & R. F. Baumeister (Eds.), *Handbook of self-regulation: Research, theory, and applications* (2nd ed., pp. 505–521). New York, NY: Guilford Press.
- Sayette, M. A., Griffin, K. M., & Sayers, W. M. (2010). Counterbalancing in smoking cue research: A critical analysis. *Nicotine & Tobacco Research*, 12, 1068–1079. doi:10.1093/ntr/ntq159
- Sayette, M. A., Loewenstein, G., Griffin, K. M., & Black, J. J. (2008). Exploring the cold-to-hot empathy gap in smokers. *Psychological Science*, 19, 926–932. doi:10.1111/j.1467-9280.2008.02178.x
- Sayette, M. A., Martin, C. S., Hull, J. G., Wertz, J. M., & Perrott, M. A. (2003). Effects of nicotine deprivation on craving response covariation in smokers. *Journal of Abnormal Psychology*, 112, 110–118. doi:10.1037/0021-843X.112.1.110
- Sayette, M. A., & Parrott, D. J. (1999). Effects of olfactory stimuli on urge reduction in smokers. *Experimental and Clinical Psychopharmacology*, 7, 151–159. doi:10.1037/1064-1297.7.2.151
- Sayette, M. A., Shiffman, S., Tiffany, S. T., Niaura, R. S., Martin, C. S., & Shadel, W. G. (2000). The measurement of drug craving. *Addiction*, 95, 189–210. doi:10.1080/09652140050111762
- Scheier, M. F., & Carver, C. S. (1985). The Self-Consciousness scale: A revised version for use with general populations. *Journal of Applied Social Psychology*, 15, 687–699. doi:10.1111/j.1559-1816.1985.tb02268.x
- Shiffman, S., Paty, J. A., Gnys, M., Kassel, J. A., & Hickcox, M. (1996). First lapses to smoking: Within-subjects analysis of real-time reports. *Journal of Consulting and Clinical Psychology*, 64, 366–379. doi:10.1037/0022-006X.64.2.366

- Shiffman, S., Paty, J. A., Kassel, J. D., Gnys, M., & Zettler-Segal, M. (1994). Smoking behavior and smoking history of tobacco chippers. *Experimental and Clinical Psychopharmacology*, *2*, 126–142. doi:10.1037/1064-1297.2.2.126
- Shiffman, S., Waters, A. J., & Hickcox, M. (2004). The Nicotine Dependence Syndrome scale: A multidimensional measure of nicotine dependence. *Nicotine & Tobacco Research*, *6*, 327–348. doi:10.1080/1462220042000202481
- Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E., Johansen-Berg, H., . . . Matthews, P. M. (2004). Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage*, *23*, S208–S219. doi:10.1016/j.neuroimage.2004.07.051
- Smolka, M. N., Buhler, M., Klein, S., Zimmermann, U., Mann, K., Heinz, A., & Braus, D. F. (2006). Severity of nicotine dependence modulates cue-induced brain activity in regions involved in motor preparation and imagery. *Psychopharmacology*, *184*, 577–588. doi:10.1007/s00213-005-0080-x
- Stippekohl, B., Winkler, M., Mucha, R. F., Pauli, P., Walter, B., Vaitl, D., & Stark, R. (2010). Neural responses to BEGIN- and END-stimuli of the smoking ritual in nonsmokers, nondeprived smokers, and deprived smokers. *Neuropsychopharmacology*, *35*, 1209–1225. doi:10.1038/npp.2009.227
- Symons, C. S., & Johnson, B. T. (1997). The self-reference effect in memory: A meta-analysis. *Psychological Bulletin*, *121*, 371–394. doi:10.1037/0033-2909.121.3.371
- Tangney, J. P., Baumeister, R. F., & Boone, A. L. (2004). High self-control predicts good adjustment, less pathology, better grades, and interpersonal success. *Journal of Personality*, *72*, 271–322. doi:10.1111/j.0022-3506.2004.00263.x
- Volkow, N. D., Fowler, J. S., Wang, G. J., Telang, F., Logan, J., Jayne, M., . . . Swanson, J. M. (2010). Cognitive control of drug craving inhibits brain reward regions in cocaine abusers. *Neuroimage*, *49*, 2536–2543. doi:10.1016/j.neuroimage.2009.10.088
- Wagner, D. D., Dal Cin, S., Sargent, J. D., Kelley, W. M., & Heatherton, T. F. (2011). Spontaneous action representation in smokers when watching movie characters smoke. *The Journal of Neuroscience*, *31*, 894–898. doi:10.1523/JNEUROSCI.5174-10.2011
- Watson, D., Clark, L. A., & Tellegen, A. (1988). Development and validation of brief measures of positive and negative affect: The PANAS scales. *Journal of Personality and Social Psychology*, *54*, 1063–1070. doi:10.1037/0022-3514.54.6.1063
- Wertz, J. M., & Sayette, M. A. (2001). A review of the effects of perceived drug use opportunity on self-reported urge. *Experimental and Clinical Psychopharmacology*, *9*, 3–13. doi:10.1037/1064-1297.9.1.3
- Wilson, S. J., Sayette, M. A., Delgado, M. R., & Fiez, J. A. (2005). Instructed smoking expectancy modulates cue-elicited neural activity: A preliminary study. *Nicotine & Tobacco Research*, *7*, 637–645. doi:10.1080/14622200500185520
- Wilson, S. J., Sayette, M. A., & Fiez, J. A. (2004). Prefrontal responses to drug cues: A neurocognitive analysis. *Nature Neuroscience*, *7*, 211–214. doi:10.1038/nn1200
- Wilson, S. J., Sayette, M. A., & Fiez, J. A. (in press). Quitting-unmotivated and quitting-motivated cigarette smokers exhibit different patterns of cue-elicited brain activation when anticipating an opportunity to smoke. *The Journal of Abnormal Psychology*.
- Woods, R. P., Cherry, S., & Mazziotta, J. (1992). Rapid automated algorithm for aligning and reslicing PET images. *Journal of Computer Assisted Tomography*, *16*, 620–633. doi:10.1097/00004728-199207000-00024
- Yalachkov, Y., Kaiser, J., & Naumer, M. J. (2009). Brain regions related to tool use and action knowledge reflect nicotine dependence. *The Journal of Neuroscience*, *29*, 4922–4929. doi:10.1523/JNEUROSCI.4891-08.2009

Received September 29, 2011

Revision received December 2, 2011

Accepted December 7, 2011 ■

E-Mail Notification of Your Latest Issue Online!

Would you like to know when the next issue of your favorite APA journal will be available online? This service is now available to you. Sign up at <http://notify.apa.org/> and you will be notified by e-mail when issues of interest to you become available!